

University of Groningen

## Value of neurohormonal and autonomic parameters for the assessment of the severity and prognosis in chronic heart failure

Tjeerdsma, Geert

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*

Publisher's PDF, also known as Version of record

*Publication date:*  
2001

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Tjeerdsma, G. (2001). *Value of neurohormonal and autonomic parameters for the assessment of the severity and prognosis in chronic heart failure*. s.n.

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

## **PART 1**

# **Introduction**

## INTRODUCTION

The phrase of “chronic heart failure” is often used and this suggests that it is well understood. However, controversy has always surrounded the use of this phrase. This is partly caused by advances in the understand of the pathophysiology, which have rendered insufficient earlier definitions. Furthermore, no single definition suffices because the clinical and physiological criteria differ. A practical definition is that CHF is “a clinical syndrome caused by an abnormality of the heart and recognized by a characteristic pattern of hemodynamic, renal, neural and hormonal responses.”<sup>1</sup> This definition implies that there should be an abnormality of the heart and that much of the clinical presentation is a consequence of the response of the body to the abnormal function of the heart.

The most common cause of CHF is no longer hypertension or valvular heart disease, as they were in past decades, but rather coronary artery disease.<sup>2</sup> The long-term existence of specific risk factors and the presence of CAD will eventually lead to the development of CHF, as demonstrated in figure 1 (modified Braunwald circle). Despite significant progress in the prevention and treatment of cardiovascular disease in the past decades, the incidence and prevalence of CHF have been increasing steadily in recent years.<sup>3</sup> Moreover, the prevalence and incidence of CHF is age dependent (figure 2):<sup>4,5-7</sup> the incidence increases from approximately 1% in persons in their fifties to over 30 % in persons older than 80 years. In normal individuals aging produces profound effects on the cardiovascular system. On an anatomic basis (figure 3), the most consistent cardiac change with aging is progressive left ventricular hypertrophy.<sup>8,9</sup> This can be exacerbated by chronic hypertension and may eventually lead to CHF.

The prognosis of patients with CHF is poor with a mortality rate 6-7 times that of the general population. The mortality one year following the onset of CHF lies around 30% .<sup>6</sup> There is no evidence to date that the prognosis of CHF in the community has improved despite advances in therapy over the last 4 decades. However, the effect of the widespread implementation of drugs that prolong life in patients with CHF may not yet have come evident on a population level. The observed increase in hospital admissions for CHF may seem paradoxical in view of the declining cardiovascular mortality and improvements in hypertension treatment in most countries in the developed world. The declining coronary artery disease mortality rates have been attributed mainly to a lower case fatality rate. It has also been suggested that the treatment of hypertension, as a major risk factor for developing CHF, merely postpones the onset of CHF to an older age rather than preventing it.<sup>6,10,11</sup>

Besides CHF, atrial fibrillation is also common in the elderly.<sup>12</sup> Owing to common risk factors including hypertension and left ventricular hypertrophy, both disorders are often present in the same patient. In addition, there is increasing evidence of a complex, reciprocal relation between CHF and atrial fibrillation. Thus, CHF may cause atrial fibrillation, electromechanical feedback and neurohumoral activation<sup>13</sup> playing an important mediating role. In addition, atrial fibrillation may promote CHF; in particular, when there is an uncontrolled ventricular rate, tachycardiomyopathy may develop and thereby CHF. Eventually, a vicious circle between CHF and atrial fibrillation may form, in which neurohormonal activation and subtle derangement of rate control are involved. Although atrial fibrillation is associated with significant morbidity and mortality during follow-up,<sup>14</sup> the prognostic significance of atrial fibrillation in the setting of CHF has not been fully elucidated.




Figure 1. The development of CHF. From risk factor to advanced CHF. CAD; coronary artery disease, LV; left ventricular, LVH; left ventricular hypertrophy.




Figure 2. Increased incidence of CHF with age (from Kannel and Belanger, Am Heart J 1991)




Figure 3. Cardiac changes in the elderly. The left atrial cavity enlarges, and the left ventricular cavity becomes smaller. The amount of space available for the mitral leaflets decreases with aging (from Roberts WC, Ann Intern Med 1972)

## NEUROHORMONAL AND AUTONOMIC CHANGES IN CHF

The observation that CHF can progress independently of the hemodynamic status of the patient has focussed interest on other possible mechanisms responsible for disease progression. As a result the neurohormonal hypothesis was formulated.<sup>15</sup> According to this hypothesis, CHF progresses because several endogenous neurohormonal systems, that are activated after the initial injury to the heart, exert a deleterious effect.<sup>16</sup> Such an effect may occur through either neurohormonally mediated hemodynamic deterioration<sup>17</sup> or through a direct toxic effect on the myocardium.<sup>18</sup> Both of these derangements may further increase neurohormonal activation and, over time, exacerbate the degree of secondary baroreceptor dysfunction.<sup>19</sup>

The first trial testing the neurohormonal hypothesis was the CONSENSUS study.<sup>20</sup> The aim was to study whether enalapril, a drug which interferes with the renin-angiotensin-aldosterone system, would reduce progression of CHF. The results demonstrated a reduction in both mortality and morbidity. This study was performed in patients with severe CHF, but favorable results were later on also obtained in the SOLVD study and V-HeFT II study.<sup>21,22</sup> It can be speculated that these effects were related to peripheral vasodilator effects (thus substantiating the hemodynamic concept), however, enalapril had the most profound effect in patients with the strongest neurohormonal activation.<sup>22</sup> Additional and intriguing evidence favoring the neurohormonal hypothesis emerged from clinical studies with drugs that block the activity of the sympathetic nervous system. Following the awareness that long-term sympathetic activation was deleterious for the heart<sup>23</sup> there was increasing interest in the use of  $\beta$ -blockers in the treatment of CHF. Although several small reports showed that  $\beta$ -blockers could cause acute worsening of CHF,<sup>24-26</sup> studies in Sweden in the early 1970s raised the possibility that long-term therapy with these drugs might produce hemodynamic and clinical benefits.<sup>27,28</sup> Additionally, the effects of  $\beta$ -blockers on mortality were studied. Although the effects on mortality were somewhat disappointing in the first small trials,<sup>29,30</sup> a profound positive effect was observed in large-scale CHF trials with  $\beta$ -blockers on top of ACE inhibitors on survival, progression of CHF, hospitalization and the incidence of sudden death.<sup>31-33</sup>

## NEUROHORMONAL AND AUTONOMIC PARAMETERS

Measurement of plasma neurohormones provides information about the extent of neurohormonal activation in CHF. The first parameter used was plasma norepinephrine level. Reports in the early 80s

demonstrated that high plasma norepinephrine levels precede and predict the progression of CHF and death even in the patients with asymptomatic CHF (figure 4).<sup>15,34,35</sup> A plasma norepinephrine level of 400-800 pg/ml corresponds with an increased mortality risk; a level over 800 pg/ml predicts an even higher 1-year mortality risk due to progressive CHF.<sup>15,34</sup> Later on, other studies demonstrated that besides plasma norepinephrine, increased levels of angiotensin-II, and aldosterone correlated significantly with mortality in CHF.<sup>36</sup> With the introduction of natriuretic peptides kits new possibilities have emerged.<sup>37,38</sup> Indeed, natriuretic peptides have shown to be superior to traditional neurohormones in the assessment of clinical severity and prognosis.<sup>39,40</sup> Furthermore, their reproducibility is higher,<sup>41</sup> which make them even more useful in clinical practice.<sup>42</sup> It has been shown that BNP is the most sensitive marker in this respect.<sup>39,43-45</sup>

Numerous studies have evaluated the autonomic nervous system in patients with CHF and have demonstrated a characteristic dysfunction of the sympathetic<sup>46</sup> and parasympathetic<sup>47</sup> limb and a deranged baroreceptor function.<sup>48,49</sup> Taken together, CHF is characterized by increased sympathetic drive and reduced parasympathetic tone. As a result of the increased sympathetic drive the myocytes of the failing heart lose their ability to respond to the  $\beta$ -adrenergic receptor agonists through a process of desensitization.<sup>46</sup> Desensitization is likely to be both beneficial and deleterious in patients with CHF. By reducing energy expenditure by the energy-starved myocardium, this adaptive response is beneficial; however, by reducing contractility and depressing the ability of the failing heart to increase its output in response to exercise, desensitization is also deleterious. The role of the parasympathetic system is not completely understood, but this system probably plays a role in inhibiting sympathetic outflow. In the setting of CHF this inhibitory effect is reduced.<sup>50</sup> Baroreceptors are the principal modulators of sympathetic and parasympathetic nerve activity and respond to changes in pressure (arterial baroreceptors) and volume (cardiopulmonary baroreceptors).<sup>51,52</sup> By regulating sympathetic and parasympathetic outflow from the brain stem, these baroreceptors provide an important control mechanism for vascular tone, heart rate and contractility, thus indirectly altering the activity of the renin-angiotensin-aldosterone system and stimulating the release of vasoactive hormones. Patients with CHF have diminished baroreflexes<sup>53</sup> and, as a result, a loss of normal responses to hemodynamic stress.<sup>54-58</sup> Various invasive and non-invasive techniques have been evaluated to study autonomic function. The analysis of HRV is an established non-invasive tool to assess autonomic function. Reduced HRV reflects the summation of the autonomic imbalances typically observed during CHF: excess sympathetic tone, parasympathetic withdrawal, and reduced baroreceptor sensitivity.<sup>59,60</sup> Autonomic function testing, based on cardiovascular reflexes, is

another method to assess autonomic status; the use of several well defined 'stress' tests (the so called Ewing battery <sup>61</sup>) provides information regarding sympathetic and parasympathetic function. Baroreflex function can be assessed invasively using drugs (phenylephrine [vasopressor] or nitroprusside [vasodepressor]) or non-invasively using a technique which directly measures muscle sympathetic nerve traffic (i.e. microneurography) <sup>62</sup> or a more recently introduced indirect technique which studies the relation between changes in arterial blood pressure and heart rate (i.e. baroreflex sensitivity). <sup>63</sup>




Figure 4. Predicted survival curves based on initial measurement of plasma norepinephrine (from Cohn JN, N Engl J Med 1984)

## AIMS OF THE THESIS

The first aim was to evaluate tools to monitor patients with CHF. In appendix 1, asymptomatic patients treated with cardiotoxic drugs for breast cancer were evaluated using HRV. It was hypothesized that early autonomic impairment occurs after treatment in the absence of left ventricular systolic dysfunction. In appendix 2, a new rapid natriuretic peptide assay (measuring BNP levels in whole blood) was evaluated to test the hypothesis that rapid measurement of BNP levels is comparable with conventional BNP measurement. In appendix 3, the technique of lower body negative pressure to simulate hypovolemia was used to test the hypothesis that neurohormonal and hemodynamic responses are impaired in the setting of CHF. In appendix 4, the relation between neurohormonal activation and the duration of concomitant atrial fibrillation in patients with CHF was evaluated.

The second aim was to study the progression and prognosis of CHF. In appendix 5, the concept was elaborated that plasma neurohormones are strongly associated with the prognosis in CHF. In appendix 6 the additional effect of atrial fibrillation in CHF patients was evaluated.

The third aim was to study drugs which are considered to have an effect on the autonomic and neurohormonal profile of patients with CHF. In appendix 7, the effect of mibefradil, a T-channel selective calcium channel blocker, on HRV parameters was studied in CHF. In appendix 8, autonomic function tests (Ewing battery) and HRV analysis were performed, before and during treatment with a  $\beta$ -blocker (metoprolol). Finally, in appendix 9, the effect of treatment with a selective dopa-mine agonist on plasma neurohormones, HRV and hemodynamics was evaluated.

## References

1. Poole-Wilson, PA: Heart Failure. *Med Int* 1985;2:866-871
2. Bourassa MG, Gurne O, Bangdiwala SI, et al: Natural history and patterns of current practice in heart failure. The Studies of Left Ventricular Dysfunction (SOLVD) Investigators. *J Am Coll Cardiol* 1993;22:14A-9A
3. Ghali JK, Cooper R, Ford E<sup>o</sup>: Trends in hospitalization rates for heart failure in the United States, 1973-1986. Evidence for increasing population prevalence. *Arch Intern Med* 1990;150:769-73.
4. O'Connell JB, Bristow MR: Economic impact of heart failure in the United States: time for a different approach. *J Heart Lung Transplant* 1994;13:S107-12
5. Detailed Diagnosis and Procedures, National Hospital Discharge Survey. 1990;92-1774
6. Ho KK, Anderson KM, Kannel WB, et al: Survival after the onset of congestive heart failure in Framingham Heart Study subjects. *Circulation* 1993;88:107-15.
7. Williams JF, Bristow MR, Fowler MB, et al: Guidelines for the evaluation and management of heart failure. Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on Evaluation and Management of Heart Failure). *J Am Coll Cardiol* 1995;26:1376-98
8. Gerstenblith G, Frederiksen J, Yin FC, et al: Echocardiographic assessment of a normal adult aging population. *Circulation* 1977;56:273-8.
9. Wei JY: Age and the cardiovascular system. *N Engl J Med* 1992;327:1735-9.
10. Yusuf S, Thom T, Abbott RD: Changes in hypertension treatment and in congestive heart failure mortality in the United States. *Hypertension* 1989;13:174-179.
11. Senni M, Tribouilloy CM, Rodeheffer RJ, et al: Congestive heart failure in the community: trends in incidence and survival in a 10-year period. *Arch Intern Med* 1999;159:29-34.
12. Benjamin EJ, Levy D, Vaziri SM, et al: Independent risk factors for atrial fibrillation in a population-based cohort. The Framingham Heart Study. *JAMA* 1994;271:840-4.
13. Tuinenburg AE, Van Veldhuisen DJ, Boomsma F, et al: Comparison of plasma neurohormones in congestive heart failure patients with atrial fibrillation versus patients with sinus rhythm. *Am J Cardiol* 1998;81:1207-10.
14. Benjamin EJ, Wolf PA, D'Agostino RB, et al: Impact of atrial fibrillation on the risk of death: the Framingham Heart Study [see comments]. *Circulation* 1998;98:946-52.
15. Packer M, Lee WH, Kessler PD, et al: Role of neurohormonal mechanisms in determining survival in patients with severe chronic heart failure. *Circulation* 1987;75:IV80-IV92.
16. Francis GS, Benedict C, Johnstone DE, et al: Comparison of neuroendocrine activation in patients with left ventricular dysfunction with and without congestive heart failure. A substudy of the Studies of Left Ventricular Dysfunction (SOLVD). *Circulation* 1990;82:1724-9.
17. Packer M: New concepts in the pathophysiology of heart failure: beneficial and deleterious interaction of endogenous haemodynamic and neurohormonal mechanisms. *J Intern Med* 1996;239:327-33.
18. Tan LB, Jalil JE, Pick R, et al: Cardiac myocyte necrosis induced by angiotensin II. *Circ Res* 1991;69:1185-95.
19. Hirsch AT, Dzau VJ, Creager MA: Baroreceptor function in congestive heart failure: effect on neurohumoral activation and regional vascular resistance. *Circulation* 1987;75:IV36-48.
20. The CONSENSUS Trial Study Group: Effects of enalapril on mortality in severe congestive heart failure. Results of the Cooperative North Scandinavian Enalapril Survival Study (CONSENSUS). *N Engl J Med* 1987;316:1429-35.
21. The SOLVD investigators: Effect of enalapril on survival in patients with reduced left ventricular ejection fractions and congestive heart failure. *N Engl J Med* 1991;325:293-302.
22. Cohn JN, Johnson G, Ziesche S, et al: A comparison of enalapril with hydralazine-isosorbide dinitrate in the treatment of chronic congestive heart failure. *N Engl J Med* 1991;325:303-10.
23. Thomas JA, Marks BH: Plasma norepinephrine in congestive heart failure. *Am J Cardiol* 1978;41:233-43.
24. Epstein SE, Braunwald E: The effect of beta adrenergic blockade on patterns of urinary sodium excretion. Studies in normal subjects and in patients with heart disease. *Ann Intern Med* 1966;65:20-7.
25. Currie PJ, Kelly MJ, McKenzie A, et al: Oral beta-adrenergic blockade with metoprolol in chronic severe dilated cardiomyopathy. *J Am Coll Cardiol* 1984;3:203-9.
26. Ikram H, Fitzpatrick D: Double-blind trial of chronic oral beta blockade in congestive cardiomyopathy. *Lancet* 1981;2:490-3.
27. Waagstein F, Hjalmarson A, Varnauskas E, et al: Effect of chronic beta-adrenergic receptor blockade in congestive cardiomyopathy. *Br Heart J* 1975;37:1022-36.
28. Swedberg K, Hjalmarson A, Waagstein F, et al: Prolongation of survival in congestive cardiomyopathy by beta-receptor blockade. *Lancet* 1979;1:1374-6.
29. Waagstein F, Bristow MR, Swedberg K, et al: Beneficial effects of metoprolol in idiopathic dilated cardiomyopathy. Metoprolol in Dilated Cardiomyopathy (MDC) Trial Study Group. *Lancet* 1993;342:1441-6.
30. CIBIS Investigators and Committees: A randomized trial of beta-blockade in heart failure. The Cardiac Insufficiency Bisoprolol Study (CIBIS). *Circulation* 1994;90:1765-73.
31. Packer M, Bristow MR, Cohn JN, et al: The effect of carvedilol on morbidity and mortality in patients with chronic heart failure. *N Engl J Med* 1996;334:1349-55.
32. CIBIS Investigators and Committees: The Cardiac Insufficiency Bisoprolol Study II (CIBIS-II): a randomised trial. *Lancet* 1999;353:9-13.
33. MERIT-HF study group: Effect of metoprolol CR/XL in chronic heart failure: Metoprolol CR/XL Randomised Intervention Trial in Congestive Heart Failure (MERIT-HF). *Lancet* 1999;353:2001-7.
34. Cohn JN, Levine TB, Olivari MT, et al: Plasma norepinephrine as a guide to prognosis in patients with chronic congestive heart failure. *N Engl J Med* 1984;311:819-23.
35. Francis GS, Cohn JN, Johnson G, et al: Plasma norepinephrine, plasma renin activity, and congestive heart failure. Relations to survival and the effects of therapy in V-HEFT II. *Circulation* 1993;87:VI-40-VI-48.
36. Swedberg K, Eneroth P, Kjeksus J, et al: Hormones regulating cardiovascular function in patients with severe congestive heart failure and their relation to mortality. CONSENSUS Trial Study Group. *Circulation* 1990;82:1730-6.
37. Colucci WS, Elkayam U, Horton DP, et al: Intravenous nesiritide, a natriuretic peptide, in the treatment of decompensated congestive heart failure. *N Engl J Med* 2000;343:246-53.
38. Dries DL, Stevenson LW: Brain natriuretic peptide as bridge to therapy for heart failure. *Lancet* 2000;355:1112-3.



39. Omland T, Aakvaag A, Bonarjee VV, et al: Plasma brain natriuretic peptide as an indicator of left ventricular systolic function and long-term survival after acute myocardial infarction. Comparison with plasma atrial natriuretic peptide and N-terminal proatrial natriuretic peptide. *Circulation* 1996;93:1963-9.
40. Richards AM, Doughty R, Nicholls MG, et al: Neurohumoral prediction of benefit from carvedilol in ischemic left ventricular dysfunction. Australia-New Zealand Heart Failure Group. *Circulation* 1999;99:786-92.
41. Murdoch DR, Byrne J, Morton JJ, et al: Brain natriuretic peptide is stable in whole blood and can be measured using a simple rapid assay: implications for clinical practice. *Heart* 1997;78:594-7.
42. Cheng V, Kazanagra R, Garcia A, et al: A rapid bedside test for B-type peptide predicts treatment outcomes in patients admitted for decompensated heart failure: a pilot study. *J Am Coll Cardiol* 2001;37:386-91.
43. McDonagh TA, Robb SD, Murdoch DR, et al: Biochemical detection of left-ventricular systolic dysfunction. *Lancet* 1998;351:9-13.
44. Motwani JG, McAlpine H, Kennedy N, et al: Plasma brain natriuretic peptide as an indicator for angiotensin-converting-enzyme inhibition after myocardial infarction. *Lancet* 1993;341:1109-13.
45. Troughton RW, Frampton CM, Yandle TG, et al: Treatment of heart failure guided by plasma aminoterminal brain natriuretic peptide (N-BNP) concentrations. *Lancet* 2000;355:1126-30.
46. Bristow MR, Ginsburg R, Minobe W, et al: Decreased catecholamine sensitivity and beta-adrenergic-receptor density in failing human hearts. *N Engl J Med* 1982;307:205-11.
47. Eckberg DL, Drabinsky M, Braunwald E: Defective cardiac parasympathetic control in patients with heart disease. *N Engl J Med* 1971;285:877-83.
48. Schwartz PJ, Zaza A, Pala M, et al: Baroreflex sensitivity and its evolution during the first year after myocardial infarction. *J Am Coll Cardiol* 1988;12:629-36.
49. Creager MA: Baroreceptor reflex function in congestive heart failure. *Am J Cardiol* 1992;69:10G-5G.
50. Azevedo ER, Parker JD: Parasympathetic control of cardiac sympathetic activity: normal ventricular function versus congestive heart failure. *Circulation* 1999;100:274-9.
51. Mancia G, Mark AL: Arterial baroreflexes in humans. Handbook of Physiology: Peripheral Circulation and Organ Blood Flow. Bethesda, Md: *American Physiological Society*; 1983. p. 755-94.
52. Mark AL, Mancia G: Cardiopulmonary baroreflexes in humans. In: Shepherd JT, Abboud FM, editors. Handbook of Physiology: Peripheral Circulation and Organ Blood Flow. Bethesda, MD: *American Physiology Society*; 1983. p. 795-813.
53. Ellenbogen KA, Mohanty PK, Szentpetery S, et al: Arterial baroreflex abnormalities in heart failure. Reversal after orthotopic cardiac transplantation. *Circulation* 1989;79:51-8.
54. Creager MA, Creager SJ: Arterial baroreflex regulation of blood pressure in patients with congestive heart failure. *J Am Coll Cardiol* 1994;23:401-5.
55. Grassi G, Seravalle G, Cattaneo BM, et al: Sympathetic activation and loss of reflex sympathetic control in mild congestive heart failure. *Circulation* 1995;92:3206-11.
56. Cody RJ, Franklin KW, Kluger J, et al: Mechanisms governing the postural response and baroreceptor abnormalities in chronic congestive heart failure: effects of acute and long-term converting-enzyme inhibition. *Circulation* 1982;66:135-42.
57. Stewart DJ, Cernacek P, Costello KB, Rouleau JL. Elevated endothelin-1 in heart failure and loss of normal response to postural change. *Circulation* 1992;85:510-7.
58. Ferguson DW, Berg WJ, Roach PJ, et al: Effects of heart failure on baroreflex control of sympathetic neural activity. *Am J Cardiol* 1992;69:523-31.
59. Saul JP, Arai Y, Berger RD, et al: Assessment of autonomic regulation in chronic congestive heart failure by heart rate spectral analysis. *Am J Cardiol* 1988;61:1292-9.
60. Brouwer J: Autonomic control of the heart in cardiovascular disease [dissertation]. University Hospital Groningen; 1996.
61. Rodrigues EA, Ewing DJ: Immediate heart rate response to lying down: simple test for cardiac parasympathetic damage in diabetics. *BMJ* 1983;287:800.
62. Mancia G, Grassi G: Baroreceptor control of the circulation in man. An update. *Clin Exp Hypertens* 1995;17:387-97.
63. Bertinieri G, Di RM, Cavallazzi A, et al: A new approach to analysis of the arterial baroreflex. *J Hypertens Suppl* 1985;3:S79-S81.